

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : B65D 81/32, 21/02, A61K 7/00, 7/48		A1	(11) International Publication Number: WO 96/37420
			(43) International Publication Date: 28 November 1996 (28.11.96)
(21) International Application Number: PCT/EP96/02274			(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 24 May 1996 (24.05.96)			
(30) Priority Data:			
08/451,940 26 May 1995 (26.05.95) US			
60/005,188 13 May 1996 (13.05.96) US			
(71) Applicant (for AU BB CA GB IE KE LK LS MN MW NZ SD SG SK SZ TT UG only): UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB).			
(71) Applicant (for all designated States except AU BB CA GB IE KE LK LS MN MW NZ SD SG SZ TT UG): UNILEVER N.V. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).			
(72) Inventors: SUARES, Alan, Joseph; 339 Wallingford Road, Cheshire, CT 06410 (US). NETTESHEIM, Susan; Apartment 4d, 61 West 62nd Street, New York, NY 10023 (US). INDURSKY, Michael; 104 Birchfield Drive, Fairfield, CT 06430 (US). BERTOLINI, Peter; 51 Country Ridge Drive, Shelton, CT 06484 (US).			
(74) Agent: EVANS, Jacqueline, G., V.; Unilever plc, Patent Division, Colworth House, Sharnbrook, Bedford MK44 1LQ (GB).			

Published

With international search report.


Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: TREATMENT REGIME FOR SKIN

(57) Abstract

A method for a skin treatment regime and a respective product is provided which cosmetically improves human skin. The product includes a first composition containing at least one active and functioning to impart a first benefit to the skin. A second composition is provided which includes a second different active and imparts a second benefit to the skin. The first and second compositions are stored in respective separate containers, which nevertheless are joined together for reminding a consumer to use the compositions in tandem and to facilitate in one sale all necessary elements of a suggested regime. Packaging suitable for this purposes can be a series of stackable jars releasably lockable together through a threaded screw arrangement one above the other.

see US 5,914,116



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

TREATMENT REGIME FOR SKINField of the Invention

5

The invention concerns a skin treatment regime formed with separate containers each storing a composition functioning to impart a benefit to the skin.

10

Background and Prior Art

15

Dual purpose single formulation cosmetic products are quite handy and popular. Examples of 2-in-1 commercial formulations are cleanser & moisturizer and shampoo & conditioner products. Unfortunately, single formulations often compromise the performance of the severally combined actives.

20

A response to this problem has been the development of skin treatment regimes. Two or more different cosmetic compositions are employed in these regimes and applied to the skin in sequential order. Illustrative commercial treatment regime compositions are cleansers, moisturizers, toners and facial foundations.

25

30

When a skin treatment regime requires multi-composition use, there have been problems with consumer education and discipline. Education is particularly a problem in mass market outlets where a cosmetic knowledgeable sales staff is generally absent. A customer may purchase one product in a treatment regime but may be uninformed with respect to a necessary complementary regime product. Even a somewhat educated customer may select the wrong complementary product. Most frustrating is when a totally educated customer discovers that the store either does not carry the

35

complementary product or is temporarily out of stock.

Even when a customer has been educated and supplied with proper products, the products may become separated at home. Moreover, the necessary reinforcement of use according to a regime may no longer be present.

Accordingly, it is an object of the present invention to provide a cosmetic product for a skin treatment regime that assists the consumer in maintaining the regime.

It is another object of the present invention to provide a cosmetic product with multi-compositions for a skin treatment regime that ensures the recommended compositions are all provided to the consumer in a single sale.

Still another object of the present invention is to provide a cosmetic product for a skin treatment regime that daily serves as a reminder to the consumer as to the proper utilization of component compositions.

Yet another object of the present invention is to provide a cosmetic product for a skin treatment regime that maintains each of the compositions together in a unit to avoid separation and misplacement within a consumer's home.

SUMMARY OF THE INVENTION

A method is provided for cosmetically improving human skin through a treatment regime including:

providing a first composition containing at least one first active within a first pharmaceutically acceptable carrier, the first composition functioning to impart a first benefit to the skin;

5 providing a second composition containing at least one second active within a second pharmaceutically acceptable carrier, the second composition functioning to impart a second benefit to the skin, the first and second actives and benefits being different from one another;

10 storing the first composition in a first container;
storing the second composition in a second container, the first and second containers being joined together;

15 applying the first composition to the skin to achieve the first benefit; and

applying the second composition to the skin after application of the first composition to achieve the second benefit.

20 Preferably the first and second containers are fitted together in a releasably joinable manner, most preferably fitted together with a coupling mechanism such as a screw thread arrangement. Advantageously the first and second containers are jars stacked one above the other and releasably locked together through the coupling mechanism.

25 Also provided is a skin treatment regime product including:

30 a first composition containing at least one first active within a first pharmaceutically acceptable carrier, the first composition functioning to impart a first benefit to the skin;

35

a second composition containing at least one second active within a second pharmaceutically acceptable carrier, the second composition functioning to impart a second benefit to the skin;

5

a first container for storing the first composition; and

10

a second container for storing the second composition, the first and second containers being joined together, preferably in a releasably joined manner.

BRIEF DESCRIPTION OF THE DRAWING

15

The above features, advantages and objects of the present invention will more fully be appreciated through the following detailed discussion, reference being made to the drawing in which:

20

Fig. 1 is a perspective view of a stacked pair of jars releasably lockable one on top another; and

25

Fig. 2 is a perspective view of a dual compartment package with pump mechanisms associated with each compartment for dispensing respective compositions.

DETAILED DESCRIPTION

30

Now it has been found that a skin treatment regime requiring application of a multiple of separate compositions can be sold to a consumer as a unit, serve as a reminder for joint usage and also educate the consumer in proper application thereof. These advantages are achieved through disposition and sale of the respective compositions in respective containers wherein the two containers are joined together.

35

Figure 1 illustrates a first embodiment of a dispensing package including a first container or jar 2, a second container or jar 4 and a domed cap 6. First container 2 is fitted at an open end thereof with a threaded male screw 8 that can lockingly engage cap 6 for closure of that jar. A first composition incorporating a first active is stored within container 2.

Second container 4 at an open end thereof is also fitted with a threaded male screw 10. A second composition is stored within the second container. Threaded male screw 10 is lockingly joinable with a threaded female screw 12 arranged below a closed bottom end 14 of the first container 2. In a preferred embodiment, the first container 2 storing first composition has an outer color different from that of the outer color of the second container 4. For instance, if the first and second compositions are respectively intended for day and night use, the first container 2 may be colored predominantly white (day formula) and the second container 4 may be colored predominantly black (night formula).

The term "predominantly" means greater than 50% of the surface area of outer walls of the container. Color coding provides the consumer with an educational signal for proper use of the cosmetic product.

Figure 2 illustrates a second embodiment of a dispensing package suitable for a lotion or other relatively non-viscous liquid cosmetic composition. This embodiment of the dispenser includes a first and second container 3, 5 having respective first and second top ends 7, 9 and first and second bottom ends 11, 13. The top ends are open and the bottom ends are closed. First and second pump mechanism 15, 16 with respective dispensing nozzles 18, 20 having

dispensing openings 22, 24 are positioned over the top ends of their respective containers. Openings 22, 24 of the dispensing nozzles are oriented in a direction 180° opposite one another. It is an important feature that the dispensing nozzle openings are oriented so that simultaneous pressing of both pump mechanisms is not suggested to a consumer. A manner for avoiding this orients the dispensing nozzle openings at an angle of at least 60°, most preferably 180° apart from one another.

10

Table I lists various types of cosmetic compositions which may serve as the first and second compositions referred to above. These are described in terms of their major benefit to the skin.

TABLE I

FIRST COMPOSITION	SECOND COMPOSITION
Cleanser	Moisturizer
Cleanser	Anti-acne preparation
Cleanser	Moisturizer
Cleanser	Facial Foundation
Moisturizer	Toner
Cleanser	Skin Lightener
Self-Tanner	Cleanser
Cleanser	Sunscreen
Skin Lightener	Sunscreen
Sunscreen	Anti-wrinkle Cream
Moisturizer	Sunscreen

It is to be understood that this invention is not limited to just two compartments or that these contain only two separate compositions. For instance, there may be anywhere from three to six additional compartments, each containing their respective skin treatment regime. An example would be a package with a stack of four jars similar to that of Figure 1 each respectively containing a moisturizer, cleanser, toner and facial foundation. It is to be emphasized that the assortment of compositions are intended to be used in combination with one another for a treatment regime. This would contrast with, for instance, a travel kit where a variety of cosmetics could be stored but each functioning independently such as a series of colored powders or the like.

At least one active material will be incorporated into each of the separate compositions, the active being present as the primary functional ingredient to deliver the benefit for which the composition is applied.

Sunscreen compositions will of course incorporate a sunscreen as the active material. The term "sunscreen" is used to denote ultraviolet ray-blocking compounds inhibiting absorption within the wavelength region between 290 and 420 nm. These compounds may either be organic or inorganic. The organic compounds are preferred. When the sunscreen is inorganic and serves as the sole sun protective substance, it is suitably present at levels ranging from about 5 to 30%, preferably from about 8 to 15% by weight.

Typical inorganic sunscreens include titanium dioxide, zinc oxide, iron oxide and combinations thereof. Most preferred is titanium dioxide, especially having an average particle size no higher than 700 nm, preferably no higher than 200 nm, optimally less than 35 nm.

Organic sunscreens may be classified into five groups based upon their chemical structures: para-amino benzoates; salicylates; cinnamates; benzophenones; coumarins; azoles and miscellaneous chemicals including menthyl anthralinate. Also polymeric particles may be useful such as polyethylene and polyamides. Organic sunscreen compound will typically range in amount anywhere from 0.1 to 25%, optimally from 1 to 15%, most preferably from 5 to 10% by weight.

Anti-wrinkle compositions will usually contain one or more actives such as alpha-hydroxycarboxylic acids or salts thereof, beta-hydroxycarboxylic acid or salts thereof, retinoids, ceramides and combinations thereof.

A wide variety of C₂-C₃₀ alpha-hydroxycarboxylic acids may be employed. Suitable examples of which include:

α-hydroxyethanoic acid

α-hydroxypropanoic acid

- 5 α -hydroxyhexanoic acid
 α -hydroxyoctanoic acid
 α -hydroxydecanoic acid
 α -hydroxydodecanoic acid
 α -hydroxytetradecanoic acid
 α -hydroxyhexadecanoic acid
 α -hydroxyoctadecanoic acid
 α -hydroxyeicosanoic acid
 α -hydroxydocosanoic acid
10 α -hydroxyhexacosanoic acid, and
 α -hydroxyoctacosanoic acid

15 Particularly preferred from the above list are α -hydroxyethanoic acid (commonly known as glycolic acid), α -hydroxypropanoic acid (commonly known as lactic acid) and α -hydroxyoctanoic acid (commonly known as α -hydroxycaprylic acid or HCA).

20 Suitable C_2 - C_{60} beta-hydroxycarboxylic acids or salts are usually selected from derivatives of salicylic acid, especially esters thereof.

25 For purposes of this invention, the term α -hydroxycarboxylic acids and beta-hydroxycarboxylic acids are intended to include not only the acid form but also salts thereof. Typical salts are the alkalimetal, ammonium and C_2 - C_{60} ammonium salts thereof. Particularly preferred are the sodium, potassium, triethanolammonium, polyethyleneimine and ammonium salts. Combinations of all the foregoing may be
30 present in the compositions. Amounts of the α or β -hydroxycarboxylic acid or salt thereof may suitably range from 0.01 to about 15%, preferably from 0.1 to about 12%, optimally from 0.5 to about 9% by weight.

Suitable retinoids are retinol, retinoic acid or the C₁-C₂₀ esters of retinol and retinoic acid. Illustrative ceramides are Ceramide 1, Ceramide 2, Ceramide 3 and Ceramide 6. Pseudoceramides may also be useful. Levels of retinoids and ceramides may suitably range from 0.00001 to 2%, preferably from 0.0001 to 0.1% by weight.

Surfactants are included as actives for cleanser compositions of this invention. These may be selected from nonionic, anionic, cationic or amphoteric emulsifying agents. They may suitably range in an amount from 0.1 to 30% by weight. Illustrative of the nonionic surfactants are alkoxyated compounds based upon C₈-C₂₂ fatty alcohols, C₈-C₂₂ fatty acids and sorbitan.

Skin lighteners may include such actives as kojic acid, niacinamide, hydroquinone and derivatives of these materials. Amounts may typically range from 0.001 to 10% by weight.

Self-tanning compositions will usually include an active such as dihydroxyacetone, possibly in combination with amines and/or amino acids. Amounts of the combined actives may suitably range from 1% to 30% by weight.

Anti-acne preparations will usually include an active selected from the group consisting of benzoyl peroxide, an alpha-hydroxycarboxylic acid, salicylic acid, retinoids and their derivatives. Amounts of these materials may suitably range from 0.1 to 30% by weight of the composition.

The first, second or any other compositions of the present invention may be solid or liquid, aqueous or anhydrous and opaque or transparent.

A pharmaceutically acceptable carrier is ordinarily utilized to deliver the active components of the first and second compositions. Most preferably, cosmetic compositions of this invention will be in emulsion form. By definition, an
5 emulsion is a dispersed system containing at least two immiscible liquid phases, one of which is dispersed in the form of small droplets throughout the other. Water and oil are the most common immiscible phases. An emulsion in which oil is dispersed as droplets throughout the aqueous phase is
10 termed an oil-in-water emulsion. When water is the dispersed phase and an oil is the dispersion medium, a water-in-oil emulsion exists. Contemplated within the scope of this invention are emulsions in the form of lotions and creams of both types of emulsions, those where the water phase is
15 continuous and those where the oil phase is continuous. The amount of these phases may suitably range from 99:1 to 1:99 by weight.

The term "pharmaceutically acceptable carrier" according to
20 this invention includes emollients, humectants, thickeners, silicones and water. Total amount of the carrier may suitably range from 30 to 99.9%, preferably from 50 to 90%, optimally from 70 to 85% by weight.

25 A variety of oily emollients may be employed in the compositions of this invention. These emollients may be selected from one or more of the following classes:

- 30 1. Hydrocarbon oils and waxes. Examples thereof are mineral oil, petrolatum, paraffin, ceresin, ozokerite, microcrystalline wax, polyethylene, and perhydrosqualene.
- 35 2. Triglyceride esters such as vegetable and animal fats and oils. Examples include castor oil, cocoa butter,

safflower oil, cottonseed oil, corn oil, olive oil, cod liver oil, almond oil, avocado oil, palm oil, sesame oil, squalene, and maleated soybean oil.

- 5 3. Acetoglyceride esters, such as acetylated monoglycerides.
4. Ethoxylated glycerides, such as ethoxylated glyceryl monostearate.
- 10 5. Alkyl esters of fatty acids having 10 to 22 carbon atoms. Methyl, isopropyl, and butyl esters of fatty acids are useful herein. Examples include hexyl laurate, isohexyl laurate, isohexyl palmitate, isopropyl
- 15 palmitate, decyl oleate, isodecyl oleate, hexadecyl stearate, decyl stearate, isopropyl isostearate, diisopropyl adipate, diisohexyl adipate, dihexyldecyl adipate, diisopropyl sebacate, lauryl lactate, myristyl lactate, and cetyl lactate.
- 20 6. Alkenyl esters of fatty acids having 10 to 22 carbon atoms. Examples thereof include oleyl myristate, oleyl stearate, and oleyl oleate.
- 25 7. Fatty acids having 10 to 22 carbon atoms. Suitable examples include pelargonic, lauric, myristic, palmitic, stearic, isostearic, hydroxystearic, oleic, linoleic, ricinoleic, arachidic, behenic, and erucic acids.
- 30 8. Fatty alcohols having 10 to 22 carbon atoms. Lauryl, myristyl, cetyl, hexadecyl, stearyl, isostearyl, hydroxystearyl, oleyl, ricinoleyl, behenyl, erucyl, and 2-octyl dodecanyl alcohols are examples of satisfactory fatty alcohols.

9. Fatty alcohol ethers. Ethoxylated fatty alcohols of 10 to 22 carbon atoms including the lauryl, cetyl, stearyl, isostearyl, oleyl, and cholesterol alcohols, having attached thereto from 1 to 50 ethylene oxide groups or 1 to 50 propylene oxide groups.
10. Ether-esters such as fatty acid esters of ethoxylated fatty alcohols.
11. Lanolin and derivatives. Lanolin, lanolin oil, lanolin wax, lanolin alcohols, lanolin fatty acids, isopropyl lanolate, ethoxylated lanolin, ethoxylated lanolin alcohols, ethoxylated cholesterol, propoxylated lanolin alcohols, acetylated lanolin alcohols, lanolin alcohols linoleate, lanolin alcohols ricinoleate, acetate of lanolin alcohols ricinoleate, acetate of ethoxylated alcohols-esters, hydrogenolysis of lanolin, ethoxylated hydrogenated lanolin, ethoxylated sorbitol lanolin, and liquid and semisolid lanolin absorption bases are illustrative of emollients derived from lanolin.
12. Polyhydric alcohol esters. Ethylene glycol mono and di-fatty acid esters, diethylene glycol mono- and di-fatty acid esters, polyethylene glycol (200-6000) mono- and di-fatty acid esters, propylene glycol mono- and di-fatty acid esters, polypropylene glycol 2000 mono- oleate, polypropylene glycol 2000 monostearate, ethoxylated propylene glycol monostearate, glyceryl mono- and di-fatty acid esters, polyglycerol poly-fatty esters, ethoxylated glyceryl monostearate, 1,3-butylene glycol monostearate, 1,3-butylene glycol distearate, polyoxyethylene polyol fatty acid esters, sorbitan fatty acid esters, and polyoxyethylene sorbitan fatty acid esters are satisfactory polyhydric alcohol esters.

13. Wax esters such as beeswax, spermaceti, myristyl myristate, stearyl stearate.
14. Beeswax derivatives, e.g. polyoxyethylene sorbitol beeswax. These are reaction products of beeswax with ethoxylated sorbitol of varying ethylene oxide content, forming a mixture of ether esters.
15. Vegetable waxes including carnauba and candelilla waxes.
16. Phospholipids such as lecithin and derivatives.
17. Sterols. Cholesterol, cholesterol fatty acid esters are examples thereof.
18. Amides such as fatty acid amides, ethoxylated fatty acid amides, solid fatty acid alkanolamides.

Amounts of the above listed emollients may suitably range from 0.5 to 80% by weight of the total composition. Preferably the amounts of these emollients will range from 1 to 25%, optimally between 5 and 15% by weight.

Humectants of the polyhydric alcohol-type may also be included in the compositions of this invention. The humectant aids in increasing the effectiveness of the emollient, reduces scaling, stimulates removal of built-up scale and improves skin feel. Typical polyhydric alcohols include polyalkylene glycols and more preferably alkylene polyols and their derivatives, including propylene glycol, dipropylene glycol, polypropylene glycol, polyethylene glycol and derivatives thereof, sorbitol, hydroxypropyl sorbitol, hexylene glycol, 1,3-butylene glycol, 1,2,6-hexanetriol, ethoxylated glycerol, propoxylated glycerol and mixtures thereof. For best results the humectant is preferably

glycerol. The amount of humectant may suitably range from 0.1 to 40%, preferably between 1 and 15% by weight of the composition.

5 For improved lubricity, there may also be included one or more silicone oils or fluids which may be selected from a dimethyl polysiloxane, a methylphenyl polysiloxane and an alcohol-soluble silicone glycol copolymer. Preferred
10 siloxanes include dimethyl polysiloxane (CTFA name: dimethicone), a polysiloxane end-blocked with trimethyl units and polydimethylcyclsiloxane, (CTFA name: cyclomethicone). The preferred siloxanes exhibit a viscosity from 2 to 50 centistokes at 25°C. Amounts of the silicones can range from 0.1 to 80% by weight of the compositions, preferably from
15 about 1 to 20% by weight.

Both first and second compositions of the invention can also include thickeners/viscosifiers in amounts suitably from 0.01 to 10% by weight of the composition. As known to those
20 skilled in the art, the precise amount of thickeners can vary depending upon the consistency and thickness of the composition which is desired. Exemplary thickeners are magnesium aluminum silicate (Veegum®), guar gums (such as Jaguar HP-120®), xanthan gum, sodium carboxymethyl cellulose,
25 hydroxyalkyl and alkyl celluloses, and cross-linked acrylic acid polymers such as those sold by B.F. Goodrich under the Carbopol® trademark.

Preservatives can desirably be incorporated into both the
30 first and second compositions of this invention to protect against the growth of potentially harmful microorganisms. While it is in the aqueous phase that microorganisms tend to grow, microorganisms can also reside in the oil phase. As such, preservatives which have solubility in both water and
35 oil are preferably employed in the present compositions.

Suitable traditional preservatives for compositions of this invention are alkyl esters of para-hydroxybenzoic acid. Other preservatives which have more recently come into use include hydantoin derivatives, propionate salts, and a
5 variety of quaternary ammonium compounds. Cosmetic chemists are familiar with appropriate preservatives and routinely choose them to satisfy the preservative challenge test and to provide product stability. Particularly preferred
10 preservatives are methyl paraben, imidazolidinyl urea, sodium dehydroxyacetate, propyl paraben and benzyl alcohol. The preservatives should be selected having regard for the use of the composition and possible incompatibilities between the preservatives and other ingredients in the emulsion. Preservatives are preferably employed in amounts ranging from
15 0.01% to 2% by weight of the composition.

When present, the amount of water in a composition may suitably range from 1 to 99%, preferably from 20 to 90%,
20 optimally between 40 and 70% by weight.

In a preferred aspect of the present invention, the first and second compositions will each have at least five components in common. Moreover, the five components will be present at
25 essentially identical concentrations in both the first and second compositions.

The following examples will more fully illustrate the composition embodiments of this invention. All parts, percentages and proportions referred to herein and in the
30 appended claims are by weight unless otherwise indicated.

EXAMPLE 1

In a treatment regime to prevent and correct skin damage, a first composition was employed wherein sunscreens were
5 utilized as the actives and a sun protection benefit was imparted to the skin. This composition is intended for daytime protection. As part of the regime, a corrective anti-wrinkle preparation was applied at nighttime with alpha-hydroxycarboxylic acids utilized as the actives to correct
10 damage done by sunlight. This second, anti-wrinkle composition was applied for nighttime wear.

TABLE II
SUNSCREEN (PREVENT) COMPOSITION

	COMPONENT	WEIGHT %
	Parsol MCX®	4.00
5	Stearic Acid	3.00
	Finsolv TN®	3.00
	Butylene Glycol	2.00
	Glycerin	2.00
	MYRJ 59®	2.00
10	Uvinul M-40®	2.00
	Crodamol ISNP®	2.00
	Glycerol Monostearate	1.50
	Stearyl Alcohol	1.50
	Glycolic Acid (70%)	1.40
15	Triethanolamine (99%)	1.20
	Lactic Acid	1.15
	Magnesium Aluminum Silicate	1.00
	Arlacel 60®	1.00
	Silicone 1401®	0.80
20	Disodium EDTA	0.50
	Jaguar HP-120®	0.50
	Cholesterol	0.30
	Ammonia (Aqueous 26BE)	0.30
	Fragrance	0.30
25	Tween 80®	0.30
	Polyethyleneimine	0.25
	Methylparaben	0.15
	Vitamin E Acetate	0.10
	Propylparaben	0.10
30	Hydroxycaprylic Acid	0.10
	Vitamin A Palmitate	0.10
	Water	qs

TABLE III

ANTI-WRINKLE (CORRECT) COMPOSITION

5	COMPONENT	WEIGHT %
	Glycolic Acid (70%)	5.74
	Butylene Glycol	3.00
	Hetester® FAO/Fine	3.00
	Stearic Acid	3.00
10	Finsolv TN®	2.50
	Ceraphyl 230®	2.50
	Ammonia (Aqueous 26BE)	2.20
	Glycerin	2.00
	Myrj 59®	2.00
15	Polyethyleneimine	2.00
	Stearyl Alcohol	1.50
	Glycerol Monostearate	1.50
	Triethanolamine (99%)	1.20
	Magnesium Aluminum Silicate	1.00
20	SE 700	1.00
	Silicone Fluid 10	1.00
	Arlacel 60®	1.00
	Dow Corning 1401®	0.80
	Jaguar HP-120®	0.50
25	Disodium EDTA	0.50
	Sodium Stearoyl Lactylate	0.50
	Cholesterol	0.30
	Fragrance	0.30
	Tween 80®	0.30
30	Methylparaben	0.15
	Antifoam Emulsion	0.10
	Vitamin E Acetate	0.10
	Propylparaben	0.10

Hydroxycaprylic Acid	0.10
Vitamin A Palmitate	0.10
Bisabolol	0.10
Water	qs

5

EXAMPLE 2

10 This example illustrates a combination of a cleanser and skin-lightening regime. The cleanser of Table IV is first applied to the skin. The face is then washed with water, dried and then the skin-lightening preparation of Table V was applied.

15

TABLE IV
CLEANSER

COMPONENT	WEIGHT %
Glycerin	1.50
20 Polyoxyethylene hydrogenated castor oil	1.50
Sorbitan stearate	1.00
Squalane	10.00
Dipropylene glycol	5.00
25 Genistein	0.10
Water	qs 100.00

TABLE V
SKIN-LIGHTENER

COMPONENT	WEIGHT %
Polysorbate 80	1.00
Ethyl alcohol	3.00
Polyethylene glycol-600	5.00
Citric acid	0.03
Sodium citrate	0.20
1-o-Ethyltetraacetylglucosamine	0.10
Methylparaben	0.10
Fragrance	qs
Water	qs 100.00

EXAMPLE 3

This example illustrates a combination of a moisturizer and an anti-wrinkle cream regime. First the moisturizer is applied to the skin. Thereafter the anti-wrinkle cream is used to cover the moisturizer. Tables VI and VII illustrate these compositions.

TABLE VI
MOISTURIZER

COMPONENT	WEIGHT %
Glycerin α -methyl ether	10.00
Citric acid	0.03
Sodium citrate	0.07
Polyoxyethylene oleyl ether	0.50
Ethyl alcohol	5.00
Methylparaben	0.10
Fragrance	qs
Water	qs 100.00

TABLE VIIANTI-WRINKLE CREAM

COMPONENT	WEIGHT %
Polysorbate 60	3.00
Sorbitan stearate	3.00
Cetyl alcohol	1.00
Mineral oil	8.00
Glycerin	1.00
Methylparaben	0.10
Ammonium lactate	3.70
Water	qs 100.00

EXAMPLE 4

This example illustrates a regime wherein a cleanser and a toner are applied to the face. First the cleanser is applied. After at least a five minute scrub, the toner is placed on the face. Tables VIII and IX illustrate these compositions.

TABLE VIII

CLEANSER

PHASE	INGREDIENT	WEIGHT %
5	A Deionized Water	43.03
	Allantoin	0.50
	Methylparaben	0.20
	Ethylenediaminetetraacetic acid	0.02
	Imidazolidinyl urea	0.30
10	B Propylene glycol	6.00
	Magnesium aluminum silicate	1.00
	B Sodium carboxymethyl cellulose	0.10
15	C Stearyl alcohol	1.50
	Cetyl alcohol	3.00
	Glyceryl monostearate and Polyoxyethylene (100) monostearate	2.00
	Polyoxyethylene (10) soya sterols	2.50
	Polyoxyethylene (3) myristyl ether myristate	15.00
	Squalane	15.00
	Silicone Fluid 344®	12.00
	Propylparaben	0.10
	Butylparaben	0.10
	D Deionized Water	0.50
20	Dowicil 200®	0.15

TABLE IXTONER

5

10

15

PHASE	INGREDIENT	WEIGHT %
A	Deionized water	10.00
	Carbopol 941®	0.10
B	Deionized Water	82.90
	Glycerin	2.00
	Methylparaben	0.20
	Allantoin	0.30
	Imidazolidinyl urea	0.30
C	Acetylated polyoxyethylene (10) lanolin alcohol	1.00
	Cetyl/stearyl 2- ethylhexanoate	1.00
	Propylparaben	0.10
D	Deionized water	2.00
	98% Triethanolamine	0.10

CLAIMS

1. A method for cosmetically improving human skin through a treatment regime comprising:

5

providing a first composition containing at least one first active within a first pharmaceutically acceptable carrier, the first composition functioning to impart a first benefit to the skin;

10

providing a second composition containing at least one second active within a second pharmaceutically acceptable carrier, the second composition functioning to impart a second benefit to the skin, the first and second actives and benefits being different from one another;

15

storing the first composition in a first container;

20

storing the second composition in a second container, the first and second containers being joined together;

25

applying the first composition to the skin to achieve the first benefit; and applying the second composition to the skin after application of the first composition to achieve the second benefit.

2. A method according to claim 1 wherein the first and second containers are each fitted with a means for coupling same to one another.

30

3. A method according to claim 2 wherein the means for coupling is a threaded screw.

35

4. A method according to claim 2 wherein the first and second containers are stacked one above the other and releasably locked together by the means for coupling.
- 5 5. A method according to claim 1 wherein outside walls of the first and second containers are colored differently from one another.
- 10 6. A method according to claim 5 wherein the color of the outside walls of the first container is predominantly white and the color of the outside walls of the second container is of a color predominantly other than white.
- 15 7. A method according to claim 1 wherein each of the containers has its own pump mechanism for non-simultaneously dispensing each of the respective compositions.
- 20 8. A method according to claim 1 wherein the first and second compositions provide a benefit selected from the group consisting of cleansing, moisturizing, toning, anti-wrinkling, suncreening, self-tanning foundation covering and mixtures thereof.
- 25 9. A method according to claim 1 wherein the first and second compositions independently include an active material selected from the group consisting of alpha-hydroxycarboxylic acid or salts thereof, beta-hydroxycarboxylic acids or salts thereof, sunscreens, 30 retinoids, ceramides, surfactants, self-tanners and mixtures thereof.
- 35 10. A method according to claim 1 further comprising instructing consumers by placing on the containers or packaging associated therewith instructions on use of

the first and second compositions together in a sequential manner to achieve the first and second benefits in a treatment regime.

5 11. A skin treatment regime product comprising:

10 a first composition containing at least one first active within a first pharmaceutically acceptable carrier, the first composition functioning to impart a first benefit to the skin;

 a second composition containing at least one second active within a second pharmaceutically acceptable carrier, the second composition functioning to impart a second benefit to the skin;

15 a first container for storing the first composition; and a second container for storing the second composition, the first and second containers being joined together.

Fig.1.





Fig.2.



INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 96/02274

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 B65D81/32 B65D21/02 A61K7/00 A61K7/48

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 B65D A61K A47K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	NL,A,9 301 506 (CORNELIS ELIZABETH RIJLAARSDAM) 3 April 1995	1,2,5,6, 8-11
Y	see claims 1-6,8-14 see page 3, line 16-23 see figure 3	3,4,7
Y	--- DE,A,29 04 478 (BURCKHARDT) 21 August 1980 see page 4, line 1 - page 7, line 8; claims; figures	3,4
A	--- US,A,5 409 128 (MITCHELL) 25 April 1995 see abstract; figures	2-6
Y	--- EP,A,0 318 834 (FIEDLER) 7 June 1989 see abstract; figures --- -/--	7

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

13 September 1996

Date of mailing of the international search report

20.09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

SERRANO GALARRAGA, J

INTERNATIONAL SEARCH REPORT

 Int ional Application No
 PCT/EP 96/02274

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A,5 306 486 (MCCOOK JOHN P ET AL) 26 April 1994 see claims 1,2 see examples 4-6	1,8,11
A	--- EP,A,0 501 714 (UNILEVER PLC ;UNILEVER NV (NL)) 2 September 1992 see claims 12-15 see example 5	1,8,11
A	--- WO,A,94 06405 (PERRICONE NICHOLAS V) 31 March 1994 see page 3, line 20 - page 4, line 9 see page 5, line 13-23 see claims 1-9	1,8,9,11
A	--- EP,A,0 345 082 (HAYASHIBARA BIOCHEM LAB ;SANSHO SEIYAKU KK (JP); MISHIMA YUTAKA (J) 6 December 1989 see page 30, line 15-26 see claims 1-15 see examples 4-6,8-10 see examples 14-16,18-20	1,8,9,11
A	--- DE,A,39 11 089 (HEITLAND UND PETRE INTERNATION) 25 October 1990 Abstract see column 1, line 1-15 see column 1, line 65 - column 2, line 5 see column 4, line 32-36	1,2
A	--- GB,A,2 231 782 (JORDAN VALERIE LILIAN) 28 November 1990 see the whole document	1
A	--- CH,A,585 647 (SEBAMAT-CHEMIE GMBH) 15 March 1977 see the whole document -----	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/EP 96/02274

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
NL-A-9301506	03-04-95	NONE		
DE-A-2904478	21-08-80	NONE		
US-A-5409128	25-04-95	CA-A-	2145002	07-10-95
EP-A-318834	07-06-89	DE-C-	3741086	16-03-89
		US-A-	4949874	21-08-90
US-A-5306486	26-04-94	AU-A-	6141994	26-09-94
		WO-A-	9420066	15-09-94
		ZA-A-	9401172	21-08-95
EP-A-0501714	02-09-92	US-A-	5135748	04-08-92
		US-A-	5169624	08-12-92
		AU-B-	655229	08-12-94
		AU-A-	1135692	03-09-92
		CA-A-	2061679	29-08-92
		JP-B-	7014848	22-02-95
		ZA-A-	9201521	30-08-93
WO-A-9406405	31-03-94	NONE		
EP-A-0345082	06-12-89	JP-A-	2193917	31-07-90
		JP-A-	1305025	08-12-89
DE-A-3911089	25-10-90	NONE		
GB-A-2231782	28-11-90	NONE		
CH-A-585647	15-03-77	BE-A-	821257	18-04-75

THIS PAGE BLANK (USPTO)